

# Effect of soybean cropping on floral diversity

Leonardo Amthauer Gallardo and Jens Dauber



Agriculture faces a serious challenge as species diversity in agricultural landscapes declines. Grain legumes are thought to contribute to farmland biodiversity. In a survey of the international literature we established that, with the exception of soybean, there is little information on the impact of grain legumes on floral diversity of agroecosystems. According to a quantitative analysis of available data, soybean reduces weed biomass, density, and seed production when compared to other single crops, but somewhat increased them when the cropping sequence was considered. The floral diversity parameters species richness, Shannon diversity, and evenness were unaffected by soybean cultivation.

## Background

Agroecosystems are biodiversity-depleted ecosystems. The expansion of arable land and the intensification of its use has displaced natural habitats and reduced the biodiversity of entire landscapes. Since agriculture dominates land use over most of Europe, increasing on-farm biodiversity is a challenge for policymakers, scientists and land managers. Securing and enhancing the amount of semi-natural habitats, flower strips, intercropping (polyculture), extended crop rotations, the use of perennial crops, organic farming, and the increase in the production of biodiversity-enhancing arable crops are all relevant approaches. The positive impact of perennial forage legume species on agricultural habitats is well documented. Less is known about the effects of grain legumes. The question addressed here is, what can we conclude about the effects of annual grain legumes on farmland floral diversity from the existing scientific evidence. We searched the world-wide academic literature for reports of studies that compared grain legume crops with the crops they replace with respect to the biomass, cover grade, density, evenness, frequency, species



Soybean on the field. Photograph: ZALF

richness, hierarchical richness index, seed production, relative abundance and species richness of accompanying flora. It examined the crops grown and crop management as factors that might drive the effects of growing legumes on floral diversity.

Agricultural floral diversity is affected by factors such as site history, soil type and local environment variation, and microbial communities. Furthermore, management practices, such as different planting and harvest dates of the crops, tillage, and plant protection regimes, especially their timing, are very relevant. Although legumes are generally seen as crops that support biodiversity, there is little evidence if annual grain legumes integration into European crop sequences will increase floral biodiversity positively.

**Table 1.** Number of studies examining biodiversity parameters in relation to crop species and management

Driver	B	C	D	E	F	H	HRI	P	RA	S
[n] studies per organism groups or functional groups										
Crop species	6	3	6	-	1	2	-	-	-	5
Sequence	1	-	1	-	-	1	-	-	-	1
Polyculture	4	1	1	-	-	2	-	-	-	4
Fertilisation	3	-	-	-	-	1	-	-	-	1
Tillage	1	-	5	-	-	1	-	-	-	1
Weed control	3	1	4	2	-	2	1	-	1	2
Sb - Sequence	-	-	2	1	-	1	-	1	-	2
Sb - Tillage	-	-	2	1	-	4	-	-	-	2

Sb: Seedbank, B: Biomass C: Weed cover, D: Density, E: Shannon evenness, F: Frequency, H: Shannon diversity index, He: Plant height, HRI: Hierarchical richness index, P: Production (seed), RA: Relative abundance, S: Species richness

## Evidence


It was immediately obvious from the search of the literature that there is a scarcity of peer-reviewed evidence about the effects of introducing grain legumes into cropping systems on floral diversity, especially for grain legumes other than soybean. We found 53 sources for soybean. This was followed by pea (17 sources), lupin and faba bean (each two sources). Therefore we here focused only on soybean and chose 25 sources for further analysis, containing analysable information about the effect of soybean on associated flora. The plant parameters which were studied most were weed density, biomass, species richness, and Shannon diversity. Other parameters, such as seed production, weed cover, evenness and hierarchical index, were only seldom found in the literature (Table 1). The available literature covered both emerged weeds and seedbanks in agroecosystems. Data on emerged weeds show

the respective current state of weed communities in a crop whereas the weed seedbank provides information about long-term developments. Therefore both values were treated separately in our analyses.

We encountered two issues with the evidence that made further investigation difficult. There were too few publications and the experimental settings were largely inhomogeneous. Still, we extracted the relevant data from the literature to make relative difference comparisons by standardising the results of the various publications (Formula 1). Individual trials with varied location or management within an article, as well as data from different years, were treated as replications to address the lack of repetitions for the quantitative analysis. Annual data consisted primarily of a composite measurement of replicates in parcels at various times during the growing season. Four replications were required as a minimum.

$$\text{Formula 1: Relative difference} = \left( \left( \frac{\text{Higher value (BDSOY or BDoOther)}}{\text{Lower value (BDSOY or BDoOther)}} \right) * 100 \right) - 100$$

BDsoy is the value of the chosen biodiversity parameter for soy. BDoOther is the value of the chosen biodiversity parameter for the other crop with which soy is compared. The relative difference is positive (+) when the higher value is for soybean, and negative (-) when the higher value is for the other crop than soybean.



**Figure 1.** Figure 1. Mean relative differences in parameters of floral diversity of emerged weeds between soybean and widely grown alternative crops. The bars are the standard error for each mean. A positive value indicates a positive effect of soybean.


## Results

Taking all experiments and data into account, other crops (mostly maize, sorghum, sunflower and wheat) supported more floral diversity of emerged weeds than soybean. The difference amounted on average 150% for weed biomass, 135% for cover grade, 110% for density and 125% for seed production compared to soybean (Figure 1). The difference was smallest for species richness. Only maize had a lower floral diversity than soybean.


The results for the effect of cropping sequences that include soybean on the soil seedbank are less conclusive. Sites where soybean is included in the cropping sequence had a 36% higher weed seed density than sequences with less soy. Evenness, Shannon diversity, and species richness in the seedbank differed by less than 10% between soy-free sequences and sequences

with different amounts of soy (Figure 2). Longer cropping sequences tended to support a higher floral diversity as indicated by the Shannon diversity.

All polyculture measures, cover crop use, double-cropping, or intercropping resulted in reduced biomass of emerged weeds. In intercropping systems, the partner crop to soybean seems to play a decisive role in its influence on floral diversity parameters. Overall our global data analysis showed that intercropping soybean with a crop partner increased species richness by 38% compared to single soybean (Figure 3) even though weed biomass is reduced. This dynamic was also given for other studied crops compared to their intercrop. Systems with soybean tended to have increased plant diversity where tillage was reduced. Crop protection measures resulted in more uniform plant communities, while diversity remained almost unaffected.



**Figure 2.** Mean relative differences in parameters of floral diversity of seedbank between sequences with a high proportion of soybean and sequences with a low proportion of soybean. The bars are the standard error for each mean. A positive value indicates a positive effect of soybean inclusion in the sequence.



**Figure 3.** Mean relative difference in parameters of floral diversity of emerged weeds between soybean monocrops and intercrops with soybean and a partner crop. The bars are the standard error for each mean. A positive value indicates a positive effect of soybean monoculture.

## Conclusions

The evidence about the effect of soybean production on floral diversity is weak. Crop factors such as varying crop emergence times, canopy light interception, variations between autumn and spring-sown crops, and the presence of allelopathic chemicals impact weed diversity in different crop ecosystems. Crop traits and weed management are possible reasons for the low floral diversity in soybean compared to other arable crops. Soybean can successfully suppress accompanying vegetation because of its closed canopy. It reaches full ground cover earlier than corn. Furthermore, weeds are intensively controlled in soybean independent of the weed control system used (herbicide tolerance or conventional). Soybean introduction into crop sequences improved weed seedbank density despite being weak in terms of emerged weed biomass in single crop comparisons. This may be due to the short soybean growing period and the innate difference of both weed parameters. The results presented here offered some insights into the effect of soybean within polyculture systems. Using cover crops in soybean cultivation negatively impacted floral diversity parameters. The high crop plant densities in additive intercropping inhibits weed growth.

Compared with other crops, the evidence indicates that soybean crops have less weed biomass compared with other crops in cropping systems. Diversity oriented parameters such as Shannon diversity, evenness, and species richness remained almost unaffected. Surprisingly, weed seedbank density, contrary to the observations for emerged weeds biomass in crop comparisons, was positively influenced by including soybean in the crop sequence.

We conclude that the integration of soybean in European crop sequences generally has a neutral effect on floral diversity.

## Definitions

**Cover grade:** the proportion of the land or soil surface covered by plants, usually given as fraction 0-1, or percentage.

**Density:** the number of individuals per unit of area or space.

**Evenness:** how equal the distribution of individuals of species is between samples. This is a structural parameter for comparing different communities.

**Frequency:** the number of times a species occurs in a defined area in a given time.

**Species richness:** the number of species per unit area.

**Shannon diversity index:** an index of diversity based on the number of species and number of individuals per species.



Soybean flower. Photograph: Donal Murphy-Bokern

## Sources

- Amthauer Gallardo, D. A., Everwand, G., Stichnothe, H., Dauber, J. (2020). Environmental performance of cropping systems. Legumes Translated practice guide. Available from [www.legumestranslated.eu](http://www.legumestranslated.eu)
- Anderson, R. L. (2009). A 2-year small grain interval reduces need for herbicides in no-till soybean. *Weed Technology*, 23(3), 398-403.
- Cardina, J., Sparrow, D. H., & McCoy, E. L. (1995). Analysis of spatial distribution of common lambsquarters (*Chenopodium album*) in no-till soybean (*Glycine max*). *Weed Science*, 258-268.
- De la Fuente, E. B., Suárez, S. A., Lenardis, A. E., & Poggio, S. L. (2014). Intercropping sunflower and soybean in intensive farming systems: evaluating yield advantage and effect on weed and insect assemblages. *NJAS-Wageningen Journal of Life Sciences*, 70, 47-52.
- Ellsbury, M. M., Powell, J. E., Forcella, F., Woodson, W. D., Clay, S. A., & Riedell, W. E. (1998). Diversity and dominant species of ground beetle assemblages (Coleoptera: Carabidae) in crop rotation and chemical input systems for the Northern Great Plains. *Annals of the Entomological Society of America*, 91(5), 619-625.
- Gomez, P., & Gurevitch, J. (1998). Weed community responses in a maize-soybean intercrop. *Applied Vegetation Science*, 1(2), 281-288.
- Harasim, E., Gawęda, D., Wesołowski, M., Kwiatkowski, C., & Gocół, M. (2016). Cover cropping influences physico-chemical soil properties under direct drilling soybean. *Acta Agriculturae Scandinavica, Section B—Soil & Plant Science*, 66(1), 85-94.
- Kegode, G. O., Forcella, F., & Clay, S. (1999). Influence of crop rotation, tillage, and management inputs on weed seed production. *Weed Science*, 175-183.
- Légère, A., Stevenson, F. C., & Ziadi, N. (2008). Contrasting responses of weed communities and crops to 12 years of tillage and fertilisation treatments. *Weed Technology*, 22(2), 309-317.
- Martin, R. C., Voldeng, H. D., & Smith, D. L. (1990). Intercropping maize and soybean for silage in a cool-temperature region: yield, protein and economic effects. *Field Crops Research*, 23(3-4), 295-310.
- Molina, G. A., Poggio, S. L., & Ghera, C. M. (2014). Epigeal arthropod communities in intensively farmed landscapes: effects of land use mosaics, neighbourhood heterogeneity, and field position. *Agriculture, Ecosystems & Environment*, 192, 135-143.
- Murphy, S. D., Clements, D. R., Belaoussoff, S., Kevan, P. G., & Swanton, C. J. (2006). Promotion of weed species diversity and reduction of weed seedbanks with conservation tillage and crop rotation. *Weed Science*, 54(1), 69-77.
- Nord, E. A., Curran, W. S., Mortensen, D. A., Mirsky, S. B., & Jones, B. P. (2011). Integrating multiple tactics for managing weeds in high residue no-till soybean. *Agronomy Journal*, 103(5), 1542-1551.
- Palmer, M. W., & Maurer, T. A. (1997). Does diversity beget diversity? A case study of crops and weeds. *Journal of Vegetation Science*, 8(2), 235-240.
- Pannacci, E., & Tei, F. (2014). Effects of mechanical and chemical methods on weed control, weed seed rain and crop yield in maize, sunflower and soybean. *Crop Protection*, 64, 51-59.
- Rauber, R. B., Demaría, M. R., Jobbágy, E. G., Arroyo, D. N., & Poggio, S. L. (2018). Weed Communities in semiarid rainfed croplands of Central Argentina: comparison between maize (*Zea mays*) and soybean (*Glycine max*) crops. *Weed Science*, 66(3), 368-378.
- Rypstra, A. L., & Marshall, S. D. (2005). Augmentation of soil detritus affects the spider community and herbivory in a soybean agroecosystem. *Entomologia Experimentalis et Applicata*, 116(3), 149-157.
- Shrestha, B. M., McConkey, B. G., Smith, W. N., Desjardins, R. L., Campbell, C. A., Grant, B. B., & Miller, P. R. (2013). Effects of crop rotation, crop type and tillage on soil organic carbon in a semiarid climate. *Canadian Journal of Soil Science*, 93(1), 137-146.

- Smith, R. G., McSwiney, C. P., Grandy, A. S., Suwanwaree, P., Snider, R. M., & Robertson, G. P. (2008). Diversity and abundance of earthworms across an agricultural land-use intensity gradient. *Soil and Tillage Research*, 100(1-2), 83-88.
- Snyder, E. M., Curran, W. S., Karsten, H. D., Malcolm, G. M., Duiker, S. W., & Hyde, J. A. (2016). Assessment of an integrated weed management system in no-till soybean and maize. *Weed Science*, 64(4), 712-726.
- Sosnoskie, L. M., Herms, C. P., & Cardina, J. (2006). Weed seedbank community composition in a 35-yr-old tillage and rotation experiment. *Weed Science*, 54(2), 263-273.
- Sosnoskie, L. M., Herms, C. P., Cardina, J., & Webster, T. M. (2009). Seedbank and emerged weed communities following adoption of glyphosate-resistant crops in a long-term tillage and rotation study. *Weed Science*, 57(3), 261-270.
- Swanton, C. J., Booth, B. D., Chandler, K., Clements, D. R., & Shrestha, A. (2006). Management in a modified no-tillage maize-soybean-wheat rotation influences weed population and community dynamics. *Weed Science*, 54(1), 47-58.
- Swanton, C. J., Shrestha, A., Clements, D. R., Booth, B. D. & Chandler, K., 2002. Evaluation of alternative weed management systems in a modified no-tillage corn-soybean-winter wheat rotation: weed densities, crop yield and economics. *Weed Science*, 50, 504-511.
- Weil, R. R., & McFadden, M. E. (1991). Fertility and weed stress effects on performance of maize/soybean intercrop. *Agronomy Journal*, 83(4), 717-721.
- Wortman, S. E., Lindquist, J. L., Haar, M. J., & Francis, C. A. (2010). Increased weed diversity, density and above-ground biomass in long-term organic crop rotations. *Renewable Agriculture and Food Systems*, 281-295.

## About this practice note and Legumes Translated

**Authors:** Leonardo Amthauer Gallardo and Jens Dauber

**Publisher:** Thünen Institute of Biodiversity

**Production:** Donau Soja

**Permalink:** [www.zenodo.org/record/6351368](http://www.zenodo.org/record/6351368)

**Copyright:** © The Authours, 2022. Reproduction and dissemination is permitted for non-commercial purposes provided the authors and source are fully acknowledged.

This practice note was prepared within the Legumes Translated project funded by the European Union through Horizon 2020, Project Grant Number 817634.

**Citation:** Amthauer Gallardo, L. and Dauber, J., 2022. Effect of soybean cropping on floral diversity. Thünen Institute of Biodiversity. Legumes Translated Practice Note 64. [www.legumestranslated.eu](http://www.legumestranslated.eu)

The content is solely the responsibility of the authors. No warranties, expressed or implied, are made with respect to the information provided. Information relating to the use of plant protection products (pesticides) must be checked against the product label or other sources of product registration information.



This project is funded  
by the European Union

